Patent and Trademark Reduced to a 1995, no persons are required to respond to a color of the province a valid CMS control number.

REQUEST FOR ACCESS OF ABANDONED APPLICATION UNDER 37 CFR 1.14(a) In re Application of ulen et al Application Number Filed 310, 252 Examiner Group Art Unit Paper No Assistant Commissioner for Patents Washington, DC 20231 I hereby request access under 37 CFR 1.14(a)(3)(iv) to the application file record of the aboveidentified ABANDONED application, which is: (CHECK ONE) (A) referred to in United States Patent Number 5530/01 column (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No. ____ _____, filed ______ _, on page _____ of paper number (C) an application that claims the benefit of the filing date of an application that is ppen to public inspection, i.e., Application No. 07/3/0, 252, filed 2//3/89, or (D) an application in which the applicant has filed an authorization to lay open the complete application to the public. Please direct any correspondence concerning this request to the following address: Signature FOR PTO_USE Typed or printed name (initials) Unit:

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.





United States Patent [19]

Queen et al.

[11] Patent Number:

5,530,101

[45] Date of Patent:

Jun. 25, 1996

[54] HUMANIZED IMMUNOGLOBULINS

- [75] Inventors: Cary L. Queen, Los Altos; Harold E.
 - Selick, Belmont, both of Calif.
- [73] Assignee: Protein Design Labs, Inc., Mountain
 - View, Calif.
- [21] Appl. No.: 634,278
- [22] Filed: Dec. 19, 1990

Related U.S. Application Data

- [63] Continuation-in-part of Ser. No. 590,274, Sep. 28, 1990, abandoned, and a continuation-in-part of Ser. No. 310,252, Feb. 13, 1989, abandoned, which is a continuation-in-part of Ser. No. 290,975, Dec. 28, 1988, abandoned.

[56] References Cited

U.S. PATENT DOCUMENTS

4.816.397	3/1989	Boss et al 435/68
4,816,567	3/1989	Cabilly et al 530/387
4,867,973	9/1989	Geers et al
5,225,539	7/1993	Winter .

FOREIGN PATENT DOCUMENTS

0171496	2/1986	European Pat. Off
0173494	3/1986	European Pat. Off
0184187	6/1986	European Pat. Off
0239400	9/1987	European Pat. Off
0266663	6/1988	European Pat. Off
2188941	10/1987	United Kingdom .
WO86/05513	9/1986	WIPO .
WO87/02671	5/1987	WIPO .
WO89/01783	3/1989	WIPO.

OTHER PUBLICATIONS

Vitteta et al., "Redesigning Nature's Poisons to Create Anti-Tumor Reagents," Science 238:1098-1104 (1987). Ellison et al., "The nucleotide sequence of a human immunoglobulin C(gamma)₁ gene", Nucleic Acids Res. 10:4071-(1982).

Hieter et al., "Cloned Human and Mouse Kappa Immunoglobulin Constatn and J Region Genes Conserve homology in Functional Segments", Cell 22:197-207 (1980).

Sharon et al., "Expression of a $V_H C_K$ chimaeric protein in mouse myeloma cells", *Nature* 309:364-367 (1984).

Takeda et al., "Construction of chimaeric processed immunoglobulin genes containing mouse variable and human constant region sequences", *Nature* 314:452–454 (1985).

Tan et al., "A Human-Mouse Chimeric Immunoglobulin Gene with a Human Variable Region is Expressed in Mouse Myeloma Cells", J. Immunol. 135:3564-3567 (1985).

Morrison et al., "Chimeric human antibody molecules: Mouse antigen-binding domains with human constant region domains," *Proc. Natl. Acad. Sci.* 81:6851-6859 (1984).

Boulianne et al., "Production of functional chimeric mouse/ human antibody," Nature 312:643-646 (1984). Neuberger et al., "A hapten-specific chimeric IgE antibody with human physiological effector function," Nature 314:268-270 (1985).

Morrison, S. L., "Transfectomas Provide Novel Chimeric Antibodies," *Science* 229:1202–1207 (1985).

Sahagan et al., "A Genetically Engineered Murine/Human Chimeric Antibody Retains Specificity for Human Tumor-Associated Antigen", J. Immunol., 137:1066-1074 (1986). Liu et al., "Expression of mouse::human immunoglobulin heavy-chain cDNA in lymphoid cells", Gene 54:33-40 (1987).

Better et al., "Escherichia coli Secretion of an Active Chimeric Antibody Fragment", Science 240:1041-1043 (1988).

Waldmann, T. A., "The Structure, Function, and Expression of Interleukin-2 Receptors on Normal and Malignant Lymphocytes," *Science* 232:727-732 (1986).

Leonard et al., "The human receptor for T-cell growth factor," J. Biol. Chem. 260:1872-1880 (1985).

Farrar, J., "The biochemistry, biology, and role of interleukin-2 in the induction of cytotoxic T cell and antibodyforming B cell receptors," *Immunol. Rev.* 63:129-166 (1982).

Greene et al., "Growth of Human T Lymphocytes: An Analysis of Interleukin 2 and Its Cellular receptor", in *Progress in Hematology XIV*, E. Brown ed., Grune and Statton, New York (1986) pp. 283–301.

Verhoyen et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity", Science 239:1534–1536 (1988). Jones et al., "Replacing the complementarity-determining regions in a human antibody with those from a mouse", Nature 321:522-525 (1986).

Hale et al., "Remission Induction in Non-Hodgkin Lymphoma with Reshaped Human Monoclonal Antibody CAMPATH-1H", *Lancet* Dec. 17, 1988, pp. 1394-1399. Chothia, C. and A. M. Lesk, "Canonical Structures for the Hypervariable Regions of Immunoglobulins", *J. Mol. Biol.* 196:901-917 (1987).

(List continued on next page.)

Primary Examiner—Lila Feisee Attorney, Agent, or Firm—Townsend and Townsend and Crew

[57] ABSTRACT

Novel methods for producing, and compositions of, humanized immunoglobulins having one or more complementarity determining regions (CDR's) and possible additional amino acids from a donor immunoglobulin and a framework region from an accepting human immunoglobulin are provided. Each humanized immunoglobulin chain will usually comprise, in addition to the CDR's, amino acids from the donor immunoglobulin framework that are, e.g., capable of interacting with the CDR's to effect binding affinity, such as one or more amino acids which are immediately adjacent to a CDR in the donor immunoglobulin or those within about 3 A as predicted by molecular modeling. The heavy and light chains may each be designed by using any one or all of various position criteria. When combined into an intact antibody, the humanized immunoglobulins of the present invention will be substantially non-immunogenic in humans and retain substantially the same affinity as the donor immunoglobulin to the antigen, such as a protein or other compound containing an epitope.

13 Claims, 55 Drawing Sheets